




GENERALIZED LINEAR MODEL (GLM)

► Nelder and Wedderburn 1972

Problem: Target variable given feat. not always normally dist. \rightsquigarrow LM not suitable


- Target is binary (e.g., disease classification)
 \rightsquigarrow Bernoulli / Binomial distribution
- Target is count variable (e.g., number of sold products)
 \rightsquigarrow Poisson distribution
- Time until an event occurs (e.g., time until death)
 \rightsquigarrow Gamma distribution





Problem: Target variable given feat. not always normally dist. \rightsquigarrow LM not suitable


- Target is binary (e.g., disease classification)
 \rightsquigarrow Bernoulli / Binomial distribution
- Target is count variable (e.g., number of sold products)
 \rightsquigarrow Poisson distribution
- Time until an event occurs (e.g., time until death)
 \rightsquigarrow Gamma distribution



Solution: GLMs - extend LMs by allowing other distributions from exponential family


$$g(\mathbb{E}(y | \mathbf{x})) \equiv \mathbf{x}^T \theta \Leftrightarrow \mathbb{E}(y | \mathbf{x}) \equiv g^{-1}(\mathbf{x}^T \theta)$$

- Link function g links linear predictor $\mathbf{x}^T \theta$ to expectation \mathbb{E} of specified distribution of $y | \mathbf{x}$
- Link function g links linear predictor $\mathbf{x}^T \theta$ to expectation of distribution of $y | \mathbf{x}$
 \rightsquigarrow LM is special case: Gaussian distribution for $y | \mathbf{x}$ with g as identity function
- Link function g and distribution need to be specified
- High-order and interaction effects can be manually added as in LMs
- Note: Interpretation of weights depend on link function and distribution



Problem: Target variable given feat. not always normally dist. \rightarrow LM not suitable


- Logistic regression $\hat{=}$ GLM with Bernoulli distribution and logit link function:
 - Target is binary (e.g., disease classification)
 - \rightsquigarrow Bernoulli / Binomial distribution
 - Target is count variable $g(x) = \log\left(\frac{x}{1-x}\right) \Rightarrow g^{-1}(x) = \frac{1}{1 + \exp(-x)}$ (e.g., number of sold products)
 - Models probabilities for binary classification by Poisson distribution
 - Time until an event occurs (e.g., time until death)
 - \rightsquigarrow Gamma distribution




Solution: GLMs - extend LMs by allowing other distributions from exponential family

$$g(\mathbb{E}(y | \mathbf{x})) = \mathbf{x}^T \theta \Leftrightarrow \mathbb{E}(y | \mathbf{x}) = g^{-1}(\mathbf{x}^T \theta)$$


- Link function g links linear predictor $\mathbf{x}^T \theta$ to expectation \mathbb{E} of specified distribution of $y | \mathbf{x}$
 - \rightsquigarrow LM is special case: Gaussian distribution for $y | \mathbf{x}$ with g as identity function
- Link function g and distribution need to be specified





Problem: Target variable given feat. not always normally dist. \rightsquigarrow LM not suitable

- Typically, we set the threshold to 0.5 to predict classes, e.g.,
- Target is binary (e.g., disease classification)
 - Class 1 if $\pi(\mathbf{x}) > 0.5$
 - \rightsquigarrow Bernoulli / Binomial distribution
 - Class 0 if $\pi(\mathbf{x}) \leq 0.5$
- Target is count variable (e.g., number of sold products)
 - \rightsquigarrow Poisson distribution
- Time until an event occurs (e.g., time until death)
 - \rightsquigarrow Gamma distribution



Solution: GLMs - extend LMs by allowing other distributions from exponential family

$$g(\mathbb{E}(y | \mathbf{x})) = \mathbf{x}^T \boldsymbol{\theta} \Leftrightarrow \mathbb{E}(y | \mathbf{x}) = g^{-1}(\mathbf{x}^T \boldsymbol{\theta})$$

- Link function g links linear predictor $\mathbf{x}^T \boldsymbol{\theta}$ to expectation \mathbb{E} of specified distribution of $y | \mathbf{x}$
 - \rightsquigarrow LM is special case: Gaussian distribution for $y | \mathbf{x}$ with g as identity function
- Link function g and distribution need to be specified
- High-order and interaction effects can be manually added as in LMs


GENERALIZED LINEAR MODEL - (INTERPRETATION) 1972



Problem: Target variable given feat. not always normally dist. \rightsquigarrow LM not suitable

- Recall: Odds is ratio of two probabilities, odds ratio compares ratio of two odds
- Target is binary (e.g., disease classification)
 - \rightsquigarrow Bernoulli / Binomial distribution
 - \rightsquigarrow difficult to comprehend
- Target is count variable

(e.g., number of sold products)
 \rightsquigarrow Poisson distribution



● Time until an event occurs (e.g., time until death)
 \rightsquigarrow Gamma distribution

Interpretation:
Changing x_i by one unit, changes log-odds of class 1 compared to class 0 by θ_j

Solution: GLMs - extend LMs by allowing other distributions from exponential family

$$g(\mathbb{E}(y | \mathbf{x})) = \mathbf{x}^T \boldsymbol{\theta} \Leftrightarrow \mathbb{E}(y | \mathbf{x}) = g^{-1}(\mathbf{x}^T \boldsymbol{\theta})$$

- Link function g links linear predictor $\mathbf{x}^T \boldsymbol{\theta}$ to expectation \mathbb{E} of specified distribution of $y | \mathbf{x}$
 - \rightsquigarrow LM is special case: Gaussian distribution for $y | \mathbf{x}$ with g as identity function
- Link function g and distribution need to be specified
- High-order and interaction effects can be manually added as in LMs
- Note: Interpretation of weights depend on link function and distribution

GLM - LOGISTIC REGRESSION - INTERPRETATION



- **Recall:** Odds is ratio of two probabilities, odds ratio compares ratio of two odds
- Weights θ_j are interpreted linear as in LM (but w.r.t. log-odds)
- difficult to comprehend

$$\log\text{-odds} = \log\left(\frac{\pi(\mathbf{x})}{1 - \pi(\mathbf{x})}\right) = \log\left(\frac{P(y = 1)}{P(y = 0)}\right) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

Interpretation:


Changing x_j by one unit, changes log-odds of class 1 compared to class 0 by θ_j

- Odds for class 1 vs. class 0: $\text{odds} = \frac{\pi(\mathbf{x})}{1 - \pi(\mathbf{x})} = \exp(\theta_0 + \theta_1 x_1 + \dots + \theta_p x_p)$
- Instead of interpreting changes w.r.t. (log-odds), odds ratio is more common

$$= \frac{\text{odds}_{x_j+1}}{\text{odds}} = \frac{\exp(\theta_0 + \theta_1 x_1 + \dots + \theta_j (x_j + 1) + \dots + \theta_p x_p)}{\exp(\theta_0 + \theta_1 x_1 + \dots + \theta_j x_j + \dots + \theta_p x_p)} = \exp(\theta_j)$$



Interpretation: Changing x_j by one unit, changes the **odds ratio** for class 1 (compared to class 0) by the **factor** $\exp(\theta_j)$

GLM - LOGISTIC REGRESSION - EXAMPLE




- Create a binary target variable for bike rental data classes, e.g.,
 - Class 1: if "high number of bike rentals" > 70% quantile (i.e., cnt > 5531)
 - Class 0: if "low to medium number of bike rentals" (i.e., cnt ≤ 5531)
- Fit a logistic regression model (GLM with Bernoulli distribution and logit link)

	Weights	SE	p-value
(Intercept)	-8.52	1.21	0.00
seasonSPRING	1.74	0.60	0.00
seasonSUMMER	-0.86	0.77	0.26
seasonFALL	-0.64	0.55	0.25
temp	0.29	0.04	0.00
hum	-0.06	0.01	0.00
windspeed	-0.09	0.03	0.00
days_since_2011	0.02	0.00	0.00



GLM - LOGISTIC REGRESSION - EXAMPLE TATION



- Recall: Odds is quotient of two probabilities, odds ratio compares ratio of two odds
- Create a binary target variable for bike rental data:
 - Class 1: "high number of bike rentals" > 70% quantile (i.e., cnt > 5531)
 - Class 0: "low to medium number of bike rentals" (i.e., cnt ≤ 5531)
- Weights θ_j interpreted linear as in LM (but w.r.t. log-odds) \rightsquigarrow difficult to comprehend
- Fit a logistic regression model (GLM with Bernoulli distribution and logit link)

$\log\text{-odds} = \log\left(\frac{\pi(x)}{1-\pi(x)}\right) = \log\left(\frac{P(y=1)}{P(y=0)}\right) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$

	Weights	SE	p-value
(Intercept)	-8.52	1.21	0.00
seasonSPRING	1.74	0.60	0.00
seasonSUMMER	-0.86	0.77	0.26
seasonFALL	-0.64	0.55	0.25
temp	0.29	0.04	0.00
hum	-0.06	0.01	0.00
windspeed	-0.09	0.03	0.00
days_since_2011	0.02	0.00	0.00

Interpretation: Changing x_j by one unit, changes log-odds of class 1 compared to class 0 by θ_j

Interpretation

- If temp increases by 1°C , odds ratio for class 1 increases by factor $\exp(0.29) = 1.34$ compared to class 0, c.p. ($\hat{=}$ "high number of bike rentals" now 1.34 times more likely)

GLM - LOGISTIC REGRESSION - INTERPRETATION



- **Recall:** Odds is quotient of two probabilities, odds ratio compares ratio of two odds
- Weights θ_j interpreted linear as in LM (but w.r.t. log-odds) \rightsquigarrow difficult to comprehend

$$\text{log-odds} = \log\left(\frac{\pi(\mathbf{x})}{1 - \pi(\mathbf{x})}\right) = \log\left(\frac{P(y=1)}{P(y=0)}\right) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

Interpretation:

Changing x_j by one unit, changes log-odds of class 1 compared to class 0 by θ_j

- Odds for class 1 vs. class 0: $\text{odds} = \frac{\pi(\mathbf{x})}{1 - \pi(\mathbf{x})} = \exp(\theta_0 + \theta_1 x_1 + \dots + \theta_p x_p)$
- Instead of interpreting changes w.r.t. log-odds, it is more common to use *odds ratio*

$$= \frac{\text{odds}_{x_j+1}}{\text{odds}} = \frac{\exp(\theta_0 + \theta_1 x_1 + \dots + \theta_j(x_j + 1) + \dots + \theta_p x_p)}{\exp(\theta_0 + \theta_1 x_1 + \dots + \theta_j x_j + \dots + \theta_p x_p)} = \exp(\theta_j)$$

Interpretation: Changing x_j by one unit, changes the **odds ratio** for class 1 (compared to class 0) by the **factor** $\exp(\theta_j)$

GLM - LOGISTIC REGRESSION - EXAMPLE



- Create a binary target variable for bike rental data:
 - Class 1: "high number of bike rentals" $> 70\%$ quantile (i.e., $cnt > 5531$)
 - Class 0: "low to medium number of bike rentals" (i.e., $cnt \leq 5531$)
- Fit a logistic regression model (GLM with Bernoulli distribution and logit link)


	Weights	SE	p-value
(Intercept)	-8.52	1.21	0.00
seasonSPRING	1.74	0.60	0.00
seasonSUMMER	-0.86	0.77	0.26
seasonFALL	-0.64	0.55	0.25
temp	0.29	0.04	0.00
hum	-0.06	0.01	0.00
windspeed	-0.09	0.03	0.00
days_since_2011	0.02	0.00	0.00

GLM - LOGISTIC REGRESSION - EXAMPLE



- Create a binary target variable for bike rental data:
 - Class 1: "high number of bike rentals" $> 70\%$ quantile (i.e., $\text{cnt} > 5531$)
 - Class 0: "low to medium number of bike rentals" (i.e., $\text{cnt} \leq 5531$)
- Fit a logistic regression model (GLM with Bernoulli distribution and logit link)

	Weights	SE	p-value
(Intercept)	-8.52	1.21	0.00
seasonSPRING	1.74	0.60	0.00
seasonSUMMER	-0.86	0.77	0.26
seasonFALL	-0.64	0.55	0.25
temp	0.29	0.04	0.00
hum	-0.06	0.01	0.00
windspeed	-0.09	0.03	0.00
days_since_2011	0.02	0.00	0.00



Interpretation

- If temp increases by 1°C , odds ratio for class 1 increases by factor $\exp(0.29) = 1.34$ compared to class 0, c.p. ($\hat{=}$ "high number of bike rentals" now 1.34 times more likely)