Introduction to Machine Learning

Information Theory KL and Maximum Entropy





Learning goals

- Know the defining properties of the KL
- Understand the relationship between the maximum entropy principle and minimum discrimination information
- Understand the relationship between Shannon entropy and relative entropy

PROBLEMS WITH DIFFERENTIAL ENTROPY

Differential entropy compared to the Shannon entropy:

- Differential entropy can be negative
- Differential entropy is not invariant to coordinate transformations
- \Rightarrow Differential entropy is not an uncertainty measure and can not be meaningfully used in a maximum entropy framework.

In the following, we derive an alternative measure, namely the KL divergence (relative entropy), that fixes these shortcomings by taking an inductive inference viewpoint. • Caticha 2004

× 0 0 × 0 × ×

INDUCTIVE INFERENCE

We construct a "new" entropy measure S(p) just by desired properties.

Let ${\mathcal X}$ be a measurable space with $\sigma\text{-algebra}\ {\mathcal F}$ and measure μ that can be continuous or discrete.

We start with a prior distribution q over ${\mathcal X}$ dominated by μ and a constraint of the form

$$\int_{D} a(\mathbf{x}) dq(\mathbf{x}) = c \in \mathbb{R}$$

with $D \in \mathcal{F}$. The constraint function $a(\mathbf{x})$ is analogous to moment condition functions $g(\cdot)$ in the discrete case. We want to update the prior distribution q to a posterior distribution p that fulfills the constraint and is maximal w.r.t. S(p).

For this maximization to make sense, S must be transitive, i.e.,

$$\mathcal{S}(\mathcal{p}_1) < \mathcal{S}(\mathcal{p}_2), \mathcal{S}(\mathcal{p}_2) < \mathcal{S}(\mathcal{p}_3) \Rightarrow \mathcal{S}(\mathcal{p}_1) < \mathcal{S}(\mathcal{p}_3).$$

< X

CONSTRUCTING THE KL

1) Locality

The constraint must only update the prior distribution in *D*, *i.e.*, the region where it is active.



× × ×

For this, it can be shown that the non-overlapping domains of ${\cal X}$ must contribute additively to the entropy, i.e.,

$$\mathcal{S}(p) = \int \mathcal{F}(p(\mathbf{x}), \mathbf{x}) d\mu(\mathbf{x})$$

where F is an unknown function.

CONSTRUCTING THE KL / 2

2) Invariance to coordinate system



Enforcing 2) results in

$$\mathcal{S}(p) = \int \Phi\left(rac{dp}{dm}(\mathbf{x})
ight) dm(\mathbf{x})$$

where Φ is an unknown function, *m* is another measure on \mathcal{X} dominated by μ and $\frac{dp}{dm}$ the Radon–Nikodym derivative which becomes

- the quotient of the respective pmfs for discrete measures,
- the quotient of respective pdfs (if they exist) for cont. measures.

 $< \times$

CONSTRUCTING THE KL / 3

1 + 2)

 \Rightarrow *m* must be the prior distribution *q*, and our entropy measure must be understood relatively to this prior, so *S*(*p*) becomes, in fact, *S*(*p*||*q*).

3) Independent subsystems



If the prior distribution defines a subsystem of \mathcal{X} to be independent, then the priors can be independently updated, and the resulting posterior is just their product density.



CONSTRUCTING THE KL / 4

1 + 2 + 3)

Up to constants that do not change our entropy ranking, it follows that

$$S(p \| q) = -\int \log\left(rac{dp}{dq}(\mathbf{x})
ight) dp(\mathbf{x})$$

which is just the negative KL, i.e., $-D_{KL}(p||q)$.

- With our desired properties, we ended up with KL minimization
- This is called the principle of minimum discrimination information, i.e., the posterior should differ from the prior as least as possible
- This principle is meaningful for continuous and discrete RVs
- The maximum entropy principle is just a special case when \mathcal{X} is discrete and *q* is the uniform distribution.
- Analogously, Shannon entropy can always be treated as negative KL with uniform reference distribution.

