


# Introduction to Machine Learning



## Advanced Risk Minimization (Deep-Dive)

## Bias-Variance Decomposition (Deep-Dive)



### Learning goals

- Understand how to decompose the generalization error of a learner into
    - Bias of the learner
    - Variance of the learner
    - Inherent noise in the data
- Learning goals**
- Understand how to decompose the generalization error of a learner into
    - Bias of the learner
    - Variance of the learner
    - Inherent noise in the data

## BIAS-VARIANCE DECOMPOSITION

Let us take a closer look at the generalization error of a learning algorithm  $\mathcal{I}_L$ . This is the expected error of an induced model  $\hat{f}_{\mathcal{D}_n}$ , on training sets of size  $n$ , when applied to a fresh, random test observation.

$$GE_n(\mathcal{I}_L) = \mathbb{E}_{\mathcal{D}_n \sim P_{xy}^n, (\mathbf{x}, y) \sim P_{xy}} \left( L(y, \hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right) = \mathbb{E}_{\mathcal{D}_n, xy} \left( L(y, \hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right)$$

We therefore need to take the expectation over all training sets of size  $n$ , as well as the independent test observation.

We assume that the data is generated by

$$y = f_{\text{true}}(\mathbf{x}) + \epsilon,$$

with zero-mean homoskedastic error  $\epsilon \sim (0, \sigma^2)$  independent of  $\mathbf{x}$ .



## BIAS-VARIANCE DECOMPOSITION / 2

By plugging in the  $L_2$  loss  $L(y, f(\mathbf{x})) = (y - f(\mathbf{x}))^2$  we get

$$\begin{aligned} GE_n(\mathcal{I}_L) &= \mathbb{E}_{\mathcal{D}_n, xy} \left( L(y, \hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right) = \mathbb{E}_{\mathcal{D}_n, xy} \left( (y - \hat{f}_{\mathcal{D}_n}(\mathbf{x}))^2 \right) \\ &\stackrel{\text{LIE}}{=} \mathbb{E}_{xy} \left[ \underbrace{\mathbb{E}_{\mathcal{D}_n} \left( (y - \hat{f}_{\mathcal{D}_n}(\mathbf{x}))^2 \mid \mathbf{x}, y \right)}_{(*)} \right] \end{aligned}$$



Let us consider the error  $(*)$  conditioned on one fixed test observation  $(\mathbf{x}, y)$  first. (We omit the  $\mid \mathbf{x}, y$  for better readability for now.)

$$\begin{aligned} (*) &= \mathbb{E}_{\mathcal{D}_n} \left( (y - \hat{f}_{\mathcal{D}_n}(\mathbf{x}))^2 \right) \\ &= \underbrace{\mathbb{E}_{\mathcal{D}_n} (y^2)}_{=y^2} + \underbrace{\mathbb{E}_{\mathcal{D}_n} (\hat{f}_{\mathcal{D}_n}(\mathbf{x})^2)}_{(1)} - 2 \underbrace{\mathbb{E}_{\mathcal{D}_n} (y \hat{f}_{\mathcal{D}_n}(\mathbf{x}))}_{(2)} \end{aligned}$$

by using the linearity of the expectation.

## BIAS-VARIANCE DECOMPOSITION / 3



$$(*) = \mathbb{E}_{\mathcal{D}_n} \left( (y - \hat{f}_{\mathcal{D}_n}(\mathbf{x}))^2 \right) = y^2 + \underbrace{\mathbb{E}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x})^2 \right)}_{(1)} - 2 \underbrace{\mathbb{E}_{\mathcal{D}_n} \left( y \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right)}_{(2)} =$$

Using that  $\mathbb{E}(z^2) = \text{Var}(z) + \mathbb{E}^2(z)$ , we see that

$$= y^2 + \text{Var}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) + \mathbb{E}_{\mathcal{D}_n}^2 \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) - 2y \mathbb{E}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right)$$

Plug in the definition of  $y$

$$= f_{\text{true}}(\mathbf{x})^2 + 2\epsilon f_{\text{true}}(\mathbf{x}) + \epsilon^2 + \text{Var}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) + \mathbb{E}_{\mathcal{D}_n}^2 \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) - 2(f_{\text{true}}(\mathbf{x}) + \epsilon) \mathbb{E}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right)$$

Reorder terms and use the binomial formula

$$= \epsilon^2 + \text{Var}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) + \left( f_{\text{true}}(\mathbf{x}) - \mathbb{E}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) \right)^2 + 2\epsilon \left( f_{\text{true}}(\mathbf{x}) - \mathbb{E}_{\mathcal{D}_n} \left( \hat{f}_{\mathcal{D}_n}(\mathbf{x}) \right) \right)$$

## BIAS-VARIANCE DECOMPOSITION / 4



$$(*) = \epsilon^2 + \text{Var}_{\mathcal{D}_n}(\hat{f}_{\mathcal{D}_n}(\mathbf{x})) + \left( f_{\text{true}}(\mathbf{x}) - \mathbb{E}_{\mathcal{D}_n}(\hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right)^2 + 2\epsilon \left( f_{\text{true}}(\mathbf{x}) - \mathbb{E}_{\mathcal{D}_n}(\hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right)$$

Let us come back to the generalization error by taking the expectation over all fresh test observations  $(\mathbf{x}, y) \sim P_{xy}$ :

$$\begin{aligned} GE_n(\mathcal{I}_L) &= \underbrace{\sigma^2}_{\text{Variance of the data}} + \underbrace{\mathbb{E}_{xy} \left[ \text{Var}_{\mathcal{D}_n}(\hat{f}_{\mathcal{D}_n}(\mathbf{x}) \mid \mathbf{x}, y) \right]}_{\text{Variance of learner at } (\mathbf{x}, y)} \\ &+ \underbrace{\mathbb{E}_{xy} \left[ \left( f_{\text{true}}(\mathbf{x}) - \mathbb{E}_{\mathcal{D}_n}(\hat{f}_{\mathcal{D}_n}(\mathbf{x})) \right)^2 \mid \mathbf{x}, y \right]}_{\text{Squared bias of learner at } (\mathbf{x}, y)} + \underbrace{0}_{\text{As } \epsilon \text{ is zero-mean and independent}} \end{aligned}$$