


Introduction to Machine Learning



Advanced Risk Minimization

L1 Risk Minimizer (Deep-Dive)



Learning goals

- Derive the risk minimizer of the L1-loss

Learning goals

- Derive the optimal constant model for the L1-loss
- Derive the risk minimizer of the L1-loss
- Derive the optimal constant model for the L1-loss

L1-LOSS: RISK MINIMIZER / 2

* **Note** that since we are computing the derivative w.r.t. the integration boundaries, we need to use Leibniz integration rule

$$\frac{\partial}{\partial c} \left(\int_a^c g(c, y) dy \right) = g(c, c) + \int_a^c \frac{\partial}{\partial c} g(c, y) dy$$

$$\frac{\partial}{\partial c} \left(\int_c^a g(c, y) dy \right) = -g(c, c) + \int_c^a \frac{\partial}{\partial c} g(c, y) dy$$

We get

$$\begin{aligned} & \frac{\partial}{\partial c} \left(\int_{-\infty}^c -(y-c) p(y) dy + \int_c^{\infty} (y-c) p(y) dy \right) \\ &= \frac{\partial}{\partial c} \left(\int_{-\infty}^c \underbrace{-(y-c) p(y)}_{g_1(c,y)} dy \right) + \frac{\partial}{\partial c} \left(\int_c^{\infty} \underbrace{(y-c) p(y)}_{g_2(c,y)} dy \right) \\ &= \underbrace{g_1(c,c)}_{=0} + \int_{-\infty}^c \frac{\partial}{\partial c} (-(y-c)) p(y) dy - \underbrace{g_2(c,c)}_{=0} + \int_c^{\infty} \frac{\partial}{\partial c} (y-c) p(y) dy \\ &= \int_{-\infty}^c p(y) dy + \int_c^{\infty} -p(y) dy. \end{aligned}$$



L1-LOSS: OPTIMAL CONSTANT MODEL

Proof:


- Firstly note that for $n = 1$ the median $\hat{\theta} = \text{med}(y^{(i)}) = y^{(1)}$ obviously minimizes the emp. risk \mathcal{R}_{emp} using the L1 loss.
- Hence let $n > 1$ in the following For $a, b \in \mathbb{R}$, define

$$S_{a,b} : \mathbb{R} \rightarrow \mathbb{R}_0^+, \theta \mapsto |a - \theta| + |b - \theta|$$

Any $\hat{\theta} \in [a, b]$ minimizes $S_{a,b}(\theta)$, because it holds that

$$S_{a,b}(\theta) = \begin{cases} |a - b|, & \text{for } \theta \in [a, b] \\ |a - b| + 2 \cdot \min\{|a - \theta|, |b - \theta|\}, & \text{otherwise.} \end{cases}$$

$$S(\theta) = |a - \theta| + |b - \theta| \text{ for } (a,b) = (3,1)$$



L1-LOSS: OPTIMAL CONSTANT MODEL / 2

W.l.o.g. assume now that all $y^{(i)}$ are sorted in increasing order.

Let us define $i_{\max} = n/2$ for n even and $i_{\max} = (n-1)/2$ for n odd and consider the intervals

$$\mathcal{I}_i := [y^{(i)}, y^{(n+1-i)}], i \in \{1, \dots, i_{\max}\}.$$

By construction $\mathcal{I}_{j+1} \subseteq \mathcal{I}_j$ for $j \in \{1, \dots, i_{\max} - 1\}$ and $\mathcal{I}_{i_{\max}} \subseteq \mathcal{I}_i$.
With this, \mathcal{R}_{emp} can be expressed as

$$\begin{aligned} \mathcal{R}_{\text{emp}}(\theta) &= \sum_{i=1}^n L(y^{(i)}, \theta) = \sum_{i=1}^n |y^{(i)} - \theta| \\ &= \underbrace{|y^{(1)} - \theta| + |y^{(n)} - \theta|}_{=S_{y^{(1)}, y^{(n)}}(\theta)} + \underbrace{|y^{(2)} - \theta| + |y^{(n-1)} - \theta| + \dots}_{=S_{y^{(2)}, y^{(n-1)}}(\theta)} + \dots \\ &= \begin{cases} \sum_{i=1}^{i_{\max}} S_{y^{(i)}, y^{(n+1-i)}}(\theta) & \text{for } n \text{ is even} \\ \sum_{i=1}^{i_{\max}} (S_{y^{(i)}, y^{(n+1-i)}}(\theta)) + |y^{((n+1)/2)} - \theta| & \text{for } n \text{ is odd.} \end{cases} \end{aligned}$$



L1-LOSS: OPTIMAL CONSTANT MODEL / 3

From this follows that

- for “ n is even”: $\hat{\theta} \in \mathcal{I}_{i_{\max}} = [y^{(n/2)}, y^{(n/2+1)}]$ minimizes S_i for all $i \in \{1, \dots, i_{\max}\} \Rightarrow$ it minimizes \mathcal{R}_{emp} ,
- for “ n is odd”: $\hat{\theta} = y^{(n+1)/2} \in \mathcal{I}_{i_{\max}}$ minimizes S_i for all $i \in \{1, \dots, i_{\max}\}$ and it's minimal for $|y^{(n+1)/2} - \theta| \Rightarrow$ it minimizes \mathcal{R}_{emp} .

Since the median fulfills these conditions, we can conclude that it minimizes the $L1$ loss.

