

# L1-REGULARIZATION

- The L1-regularized risk of a model  $f(\mathbf{x} | \boldsymbol{\theta})$  is

$$\mathcal{R}_{\text{reg}}(\boldsymbol{\theta}) = \mathcal{R}_{\text{emp}}(\boldsymbol{\theta}) + \sum_j \lambda |\theta_j|$$

and the (sub-)gradient is:

$$\nabla_{\boldsymbol{\theta}} \mathcal{R}_{\text{emp}}(\boldsymbol{\theta}) + \lambda \text{sign}(\boldsymbol{\theta})$$

- Unlike in  $L_2$ , contribution to grad. doesn't scale with  $\theta_j$  elements.
- Again: quadratic Taylor approximation of  $\mathcal{R}_{\text{emp}}(\boldsymbol{\theta})$  around its minimizer  $\hat{\boldsymbol{\theta}}$ , then regularize:

$$\tilde{\mathcal{R}}_{\text{reg}}(\boldsymbol{\theta}) = \mathcal{R}_{\text{emp}}(\hat{\boldsymbol{\theta}}) + \frac{1}{2}(\boldsymbol{\theta} - \hat{\boldsymbol{\theta}})^T \mathbf{H}(\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}) + \sum_j \lambda |\theta_j|$$



# L1-REGULARIZATION / 2

- To cheat and simplify, we assume the  $\mathbf{H}$  is diagonal, with  $H_{j,j} \geq 0$
- Now  $\tilde{\mathcal{R}}_{\text{reg}}(\theta)$  decomposes into sum over params  $\theta_j$  (separable!):

$$\tilde{\mathcal{R}}_{\text{reg}}(\theta) = \mathcal{R}_{\text{emp}}(\hat{\theta}) + \sum_j \left[ \frac{1}{2} H_{j,j} (\theta_j - \hat{\theta}_j)^2 \right] + \sum_j \lambda |\theta_j|$$

- We can minimize analytically:

$$\begin{aligned} \hat{\theta}_{\text{lasso},j} &= \text{sign}(\hat{\theta}_j) \max \left\{ |\hat{\theta}_j| - \frac{\lambda}{H_{j,j}}, 0 \right\} \\ &= \begin{cases} \hat{\theta}_j + \frac{\lambda}{H_{j,j}} & , \text{ if } \hat{\theta}_j < -\frac{\lambda}{H_{j,j}} \\ 0 & , \text{ if } \hat{\theta}_j \in \left[ -\frac{\lambda}{H_{j,j}}, \frac{\lambda}{H_{j,j}} \right] \\ \hat{\theta}_j - \frac{\lambda}{H_{j,j}} & , \text{ if } \hat{\theta}_j > \frac{\lambda}{H_{j,j}} \end{cases} \end{aligned}$$

- Shows how lasso (approx) transforms the normal minimizer
- If  $H_{j,j} = 0$  exactly,  $\hat{\theta}_{\text{lasso},j} = 0$

